

# Open Research Online

The Open University's repository of research publications  
and other research outputs

## SMART-1 Impact Ground-based campaign

### Conference or Workshop Item

#### How to cite:

Ehrenfreund, P.; Foing, B. H.; Veillet, C.; Wooden, D.; Gurvits, L.; Cook, A. C.; Koschny, D.; Biver, N.; Buckley, D.; Ortiz, J. L.; Di Martino, M.; Dantowitz, R.; Cooke, B.; Reddy, V.; Wood, M.; Vennes, S.; Albert, L.; Sugita, S.; Kasuga, T.; Meech, K.; Tokunaga, A.; Lucey, P.; Krots, A.; Palle, E.; Montanes, P.; Trigo-Rodriguez, J.; Cremonese, G.; Barbieri, C.; Ferri, F.; Mangano, V.; Bhandari, N.; Chandrasekhar, T.; Kawano, N.; Matsumoto, K.; Taylor, C.; Hanslmeyer, A.; Vaubaillon, J.; Schultz, R.; Erd, C.; Gondoin, P.; Levasseur-Regourd, A.-C.; Khodachenko, M.; Rucker, H.; Burchell, M.; Cole, M.; Koschny, D.; Svedhem, H.; Rossi, A.; Colaprete, T.; Goldstein, D.; Schultz, P. H.; Alkalai, L.; Banerdt, B.; Kato, M.; Graham, F.; Ball, A.; Taylor, E.; Baldwin, E.; Berezhnoy, A.; Lammer, H.; Koschny, D.; Talevi, M.; Landeau-Constantin, J.; Weyhe, B. v.; Ansari, S.; Lawton, C.; Lebreton, J. P.; Friedman, L.; Betts, B.; Buoso, M.; Williams, S.; Cirou, A.; David, L.; Sanguy, O.; Burke, J. D.; Maley, P. D.; de Moraes, V. M.; Marchis, F.; Munoz, J. M. H.; Dighay, J.-L. and Taylor, C. (2007). SMART-1 Impact Ground-based campaign. In: 38th Lunar and Planetary Science Conference, 12-16 Mar 2007, League City, Texas, USA.

For guidance on citations see [FAQs](#).

© [\[not recorded\]](#)

Version: [\[not recorded\]](#)

Link(s) to article on publisher's website:

<http://www.lpi.usra.edu/meetings/lpsc2007/pdf/2446.pdf>

---

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's data [policy](#) on reuse of materials please consult the policies page.

---

## SMART-1 IMPACT GROUND-BASED CAMPAIGN

P. Ehrenfreund, B.H. Foing, C. Veillet, D. Wooden, L. Gurvits, A. C. Cook, D. Koschny, N. Biver, D. Buckley, J.L. Ortiz, M. Di Martino, R. Dantowitz, B. Cooke, V. Reddy, M. Wood, S. Vennes, L. Albert, S. Sugita, T. Kasuga, K. Meech, A. Tokunaga, P. Lucey, A. Krots, E. Palle, P. Montanes, J. Trigo-Rodriguez, G. Cremonese, C. Barbieri, F. Ferri, V. Mangano, N. Bhandari, T. Chandrasekhar, N. Kawano, K. Matsumoto, C. Taylor, A. Hanslmeyer, J. Vaubaillon, R. Schultz, C. Erd, P. Gondoin, AC Levasseur-Regourd, M. Khodachenko, H. Rucker, et al. (SMART-1 Coordinated observations group), M. Burchell, M. Cole, D. Koschny, H. Svedhem, A. Rossi, T. Colaprete, D. Goldstein, P.H. Schultz, L. Alkalai, B. Banerdt, M. Kato, F. Graham, A. Ball, E. Taylor, E. Baldwin, A. Berezhnoy, H. Lammer et al (SMART-1 impact prediction group), D. Koschny, M. Talevi, J. Landeau-Constantin, B. v. Weyhe, S. Ansari, C. Lawton, J.P. Lebreton, L. Friedman, B. Betts, M. Buoso, S. Williams, A. Cirou, L. David, O. Sanguy, J.D. Burke, P.D. Maley, V.M. de Morais, F. Marchis, J.M. H. Munoz, J.-L. Dighaye, C. Taylor et al (SMART-1 outreach and amateur astronomer coordination), *SMART1 Impact Campaign Team, ESTEC/SCI-S, postbus 299, 2200 AG Noordwijk, NL, Europe*, (P. Ehrenfreund@chem.leidenuniv.nl, Bernard.Foing@esa.int)

**Introduction:** SMART-1 was launched in 2003 and was orbiting the Moon on a 5 hours period until impact on 3 September 2006. A slight thrusting manoeuvre was operated in June to fine tune the exact time and location of impact. SMART-1 (285 kg) impacted at 5h42:22 UT at Lake of Excellence in grazing incidence with a velocity of 2 km/s and kinetic energy 600 MJ. We called the community to make predictions of impact magnitude, cloud ejecta dynamics and exospheric effects. We also organized a ground based observation campaign to perform coordinated measurements of the impact. Results from the coordinated multi-site campaign will be discussed.

## Predicted effects and proposed observations:

The science interest for SMART-1 impact observations are to study impact physics, lunar science and spacecraft related effects. The thermal and dynamic evolution of the thermal flash and ejecta right after impact can constrain the understanding of impact processes such as dust acceleration, gravity controlled excavation, and strength related effects. The monitoring of spacecraft volatiles released after impact can help to simulate and to understand processes occurring on volatile rich natural impactors. One should also monitor the speed and dynamics of ejecta and dust clouds that could be followed in the Earthshine or in sunlight, other transient changes, and the characteristics of impact crater and deposited ejecta. Among the relevant observations are infrared imaging of thermal flash and visible/infrared imaging of the dynamics of ejected clouds, measurements of motions, plumes and curtains of ejected mass, images of ejecta reaching sunlight and investigation of the mineralogy of ejecta. In particular the infrared range from 2 – 20 microns allows to constrain the present lunar minerals (pyroxene, olivine), complementary to SMART-1 diagnostics (AMIE camera colour ratio with 0.95 micron pyroxene band, and SIR 0.9 -2.5 micron spectrometry). The spectral energy distribution can provide quantitative characteristics such as mineralogical composition, porosity, crystallinity, size distribution and space weathering.

For the thermal flash bolometric magnitudes between 7–16 (during the first second) in the infrared, depending on efficiency (0.5-0.001), were predicted. In addition to thermal effects, we expected the emission from volatile elements onboard (hydrazine  $N_2H_4$  decomposing in ammonia  $NH_3$  and H compounds, detectable in Paschen and Brackett or even Balmer if the excitation temperature is sufficient). For the Hiten impact (300 kg, 2.7 km/s, 1 kg hydrazine) a flash signature detected




Fig.1 : Orbital track over the impact site imaged 10 days before with the SMART-1 AMIE camera

in the K band was ascribed to Brackett g emission. Other probe signatures may be detectable (Al melting at 600 deg C, graphite). A crater size of 5-10 m was expected.

The volume of material ejected (predicted to be  $10 \text{ m}^3$ ) will be essentially made of dust with dominant size around 15 microns (normalized per area) and the effective area of ejecta could be  $25 \text{ km}^2$ . This can lead to obscuration of the underground soil in the first minute after impact, and partial obscuration later. The ejecta can be also traced in Earth shine reflected light giving a level of  $V=14.5$  per  $\text{arcsec}^2$  accessible to large telescopes (1500 photon per second for a 2 m class telescope). Small telescopes are able to image the Earth-shine on the Moon, even with a modest resolution of 1.5 arcsec. They would be able to detect the excess brightness of the effective area of elongated ejecta with excess magnitude 13-14.

The SMART-1 impact observation campaign involved a core of participating telescopes, including: the South African Large Telescope (SALT), the Calar Alto observatory in Andalucia, Spain, the ESA Optical Ground Station (OGS) at Tenerife, Spain, the TNG telescope in La Palma, Canary Islands, Spain, the CEA Cariri observatory in Brazil, the Argentina National Telescope, the Florida Tech Robotic telescopes at Melbourne FL and Kitt Peak, MSFC lunar meteor robotic telescopes, Houston 1m, Big Bear Solar Observatory, MDM telescopes at Kitt Peak, NASA IRTF, the Canada-France-Hawaii Telescope, the Japanese Subaru Auxiliary telescopes on Hawaii, the ODIN space observatory. We acknowledge also support from Nottingham University, and the USGS.

Five radio telescopes were involved in the SMART-1 observations, coordinated by the Joint Institute for VLBI (Very Long Baseline Interferometry) in Europe (JIVE): the Medicina (INAF) 32-metre antenna in Italy, the Fortaleza (ROEN) 14-metre antenna in Brazil, the German-Chilean TIGO (BKG) 6-metre antenna in Chile, the Mount Pleasant Observatory of the University of Tasmania (Australia) and the Australia Telescope Compact Array (CSIRO).

#### References:

- Burchell and Cole (2006), see SMART-1 website  
 Foing B H, et al, SMART-1 mission to the moon: Status, first results and goals Technology and science goals; *Adv. Space Research* 37, 6-13 (2006)  
 Foing B H et al, The Science Goals of SMART-1; *Earth, Moon & Planets*, 85-86, 523-531 (2001)  
 Koschny D. and Gruen E., *Icarus* 154, 402 (2001)

Links: <http://sci.esa.int/smart-1/>,  
 SMART-1 scitech website <http://sci.esa.int/>  
 ESA public site:  
<http://www.esa.int/SPECIALS/SMART-1/index.html>




Fig.2 SMART-1 radio swan song just before impact.




Fig.3 Time series of SMART-01 impact flash and ejected debris observed from CFHT (Veillet et al)